

Ram pressure stripping in a viscous intracluster medium

Elke Roediger^{*} and Marcus Brüggen^{*}

Jacobs University Bremen, P.O. Box 750 561, 28725 Bremen, Germany

Accepted. Received; in original form

ABSTRACT

In the recent literature there is circumstantial evidence that the viscosity of the intracluster medium may not be too far from the Spitzer value. In this letter, we present two-dimensional hydrodynamical simulations of ram pressure stripping of disc galaxies in a viscous intracluster medium (ICM). The values of viscosity explored range between 0.1 and 1.0 times the Spitzer value. We find that viscosity affects the appearance and the dimensions of the galactic wakes but has very little effect on the evolution of the gas mass of the galaxy.

Key words: galaxies: spiral – galaxies: evolution – galaxies: ISM – galaxies – individual: NGC 4388 – intergalactic medium

1 INTRODUCTION

In clusters, galaxies can lose some or all of their gas by ram pressure stripping (RPS) due to their motion through the intracluster medium (ICM). Both, analytical estimates (Gunn & Gott 1972) and (hydro)dynamical simulations (e.g. Abadi et al. 1999, Quilis & Moore 2001, Schulz & Struck 2001, Vollmer et al. 2001, Marcolini et al. 2003, Roediger & Brüggen 2006, Roediger & Brüggen 2007) show that RPS can remove a significant amount of gas from galaxies, and is thus important for the evolution of galaxies and the ICM.

In addition to gas loss by ram pressure pushing, galaxies also suffer gas loss by continuous (sometimes also called turbulent or viscous) stripping (e.g. Nulsen 1982; Quilis et al. 2000; Schulz & Struck 2001; Roediger & Hensler 2005; Roediger & Brüggen 2006).

Nulsen (1982) has studied the effects of transport processes and turbulence on the flow of gas past a galaxy and has found that they could produce more stripping of gas than ram pressure alone. For turbulent stripping, he found a mass stripping rate of

$$\dot{M}_{\text{turb}} \sim \pi r^2 \rho_{\text{ICM}} v \sim 7 r_{10}^2 n_{-3} v_3 M_{\odot} \text{yr}^{-1}, \quad (1)$$

where $r = 10 r_{10}$ kpc, $\rho_{\text{ICM}} = 10^{-3} n_{-3} m_{\text{H}} \text{ g cm}^{-3}$ and $v = 1000 v_3 \text{ km s}^{-1}$. This is similar to what is found in the simulations by Roediger & Brüggen (2006, 2007) who find rates of about $\sim 1 M_{\odot} \text{yr}^{-1}$. Turbulent stripping is mainly caused by the Kelvin-Helmholtz instability, which is suppressed by viscosity that stabilises modes with wavelengths smaller than r/Re (e.g. Betchov & Criminale 1967).

^{*} E-mail: e.roediger@jacobs-university.de; m.brueggen@jacobs-university.de

There exist only few detailed studies on the wakes of stripped galaxies. Using the hydrodynamical adaptive mesh refinement code FLASH, we have studied galactic ram pressure tails in a constant ICM wind (Roediger et al. 2006) as well as for galaxies on realistic cluster orbits (Roediger & Brüggen 2008). The minimum tail width of about 20 to 30 kpc is found near the galaxy. With increasing distance to the galaxy, the tail flares to widths of 30 to 80 kpc at a distance of ~ 100 kpc behind the galaxy. Other hydrodynamical simulations using either grid codes (e.g. Quilis & Moore 2001, Marcolini et al. 2003) or smoothed particle hydrodynamics (e.g. Abadi et al. 1999, Schulz & Struck 2001) do not focus on the tails. However, according to the snapshots provided in these papers, the gas tails are similar to the ones in our simulations. Using a sticky-particle code, Vollmer et al. (e.g. 1999, 2000, 2001a, 2001b, 2003, 2004a, 2005, 2006) aim at reproducing the RPS history of individual galaxies by comparing simulations and observations. However, these simulations concentrate on the gas distribution close to the galaxy.

Recently, Oosterloo & van Gorkom (2005) presented observations of a ~ 120 kpc long and < 25 kpc wide tail of H I gas associated with NGC 4388, and also suggest that this tail is due to ram-pressure stripping of this galaxy, either in the ICM of the Virgo cluster or in the halo of the nearby elliptical galaxy M86. However, the tail of NGC 4388 is narrower than the tails in the simulations presented in Roediger & Brüggen (2006, 2008), although it seems to be flaring in a similar fashion as found in the simulation. Also the X-ray and H α tails observed by Sun et al. (2006); Sun et al. (2007); Yagi et al. (2007) are much narrower than simulated ram pressure tails: While being only ~ 7 kpc wide, they reach lengths of ~ 70 kpc and show hardly any flaring.

This difference may be caused by the microphysics of the ICM.

The viscosity of the intracluster medium has been discussed before. Based on observations of the Perseus cluster it has been suggested by Fabian et al. (2003), Fabian et al. (2003) that viscosity may play an important role in dissipating energy injected by the central AGN. Circumstantial evidence for the presence of significant ICM viscosity is also provided by an examination of the morphology of H α filaments in the Perseus cluster. Several of the filaments appear to trace well-defined arcs which argues against the presence of strong turbulence in the ICM core, possibly resulting from the action of viscosity. This idea has been tested in numerical simulations by Ruszkowski et al. (2004b,a) and Reynolds et al. (2005).

In the case of a fully ionized and unmagnetised, thermal plasma, the relevant coefficient of viscosity is given by Braginskii (1958) and Spitzer (1962) as $\mu \approx 6.0 \times 10^{-17} (\ln \Lambda / 37)^{-1} T^{5/2} \text{ g cm}^{-1} \text{ s}^{-1}$, where T is the temperature of the plasma measured in Kelvin and $\ln \Lambda$ is the Coulomb logarithm. It results from the cumulative effect of weak Coulomb collisions. The mean free path of such interactions scales with T^2 . As the sound speed scales with $T^{1/2}$, the viscosity is proportional to $T^{5/2}$.

It is customary to measure the importance of viscosity through the Reynolds number, $\text{Re} = vl/\nu$, where v and l are characteristic velocities and length-scales of the system and $\nu = \mu/\rho$ the kinematic viscosity, with ρ being the fluid density. For our case

$$\text{Re} \sim 26 v_3 r_{10} n_{-3} f_v^{-1} \left(\frac{kT}{5 \text{ keV}} \right)^{-2.5}, \quad (2)$$

which indicates that viscosity may play a role unless the viscosity is strongly suppressed, i.e. the suppression factor, f_v , is sufficiently small.

This suppression results from the cluster magnetic fields. Even weak fields lead to a tiny proton gyroradius, which results in a very efficient suppression (factor of $\sim 10^{23}$ for typical ICM conditions, Spitzer 1962) of the local viscosity perpendicular to the magnetic field. Magnetic fields in the ICM are certainly tangled or even chaotic (Clarke 2004; Enßlin et al. 2005) and will lead to a reduced macroscopic viscosity. The degree of reduction, however, is unknown. Due to the same mechanism, also thermal conduction in the ICM is suppressed perpendicular to the magnetic field lines. In a recent study, Narayan & Medvedev (2001) found an effective macroscopic thermal conductivity that is a factor of $f_v \sim 10^{-2} - 0.2$ lower than the unmagnetised value. Similar arguments may apply to the viscosity. From studies of ICM turbulence in the Coma cluster, Schuecker et al. (2004) derive an upper limit on the kinematic viscosity of the ICM of $\sim 3 \cdot 10^{29} \text{ cm}^2 \text{ s}^{-1}$. For typical ICM densities and temperatures, this corresponds to a viscosity suppression factor f_v around 0.1. Thus, we might expect the viscosity to be suppressed by some factor between 10^{-2} and unity.

The magnetic fields of the ICM (e.g. Clarke 2004; Enßlin et al. 2005) may themselves influence the appearance of ram pressure stripped galaxies. Given that the thermal pressure in the ICM dominates over the magnetic pressure, the magnetic fields should be frozen-in and follow the ICM flow. Thus, they should be generally parallel to the galaxy's

tail. Such a magnetic field structure could attenuate thermal conduction between the stripped gas and the hot ICM, and it could suppress e.g. Kelvin-Helmholtz and maybe even Rayley-Taylor instabilities in the tail. However, it is unclear how well the magnetic fields will be aligned with the ICM-ISM interfaces and how strong this suppression will be. The influence of the magnetic fields will be studied in a forthcoming paper.

Here, using adaptive-mesh, hydrodynamical simulations, we investigate the effect of a macroscopic viscosity on the stripping of gas from a galaxy.

2 METHOD

We model the ICM-ISM interaction in the galaxy's rest frame, i.e. the galaxy is exposed to an ICM flow. The work of Roediger & Brüggen (2007) showed that the classical estimate of the stripping radius given by Gunn & Gott (1972) can be adapted to galaxies exposed to a variable ICM wind. Therefore, here we decouple the effect of time-variability and use a constant ICM wind to focus on the effect of viscosity.

To include viscosity in our hydro-simulations, we add velocity diffusion to the momentum equation

$$\frac{\partial(\rho v_i)}{\partial t} + \frac{\partial}{\partial x_k}(\rho v_k v_i) + \frac{\partial P}{\partial x_i} = \rho g_i + \frac{\partial \pi_{ik}}{\partial x_k}, \quad (3)$$

with

$$\pi_{ik} = \frac{\partial}{\partial x_k} \left[2\mu \left(e_{ik} - \frac{1}{3} \Delta \delta_{ik} \right) \right], \quad (4)$$

where P is pressure, ρ density, v_i the components of velocity, g_i the components of the gravitational acceleration, δ_{ik} the Kronecker delta,

$$e_{ik} = \frac{1}{2} \left(\frac{\partial u_i}{\partial x_k} + \frac{\partial u_k}{\partial x_i} \right), \quad (5)$$

and $\Delta = e_{ii}$. Here μ is the coefficient of (shear) viscosity. In our simulations we neglected bulk viscosity since it vanishes for an ideal gas. We use the standard Spitzer viscosity for an unmagnetised plasma, for which $\mu = 6.0 \times 10^{-17} (\ln \Lambda / 37)^{-1} T^{5/2} f_v \text{ g cm}^{-1} \text{ s}^{-1}$.

We have noted earlier that the precise value of the suppression factor is highly uncertain and, depending on the nature of magnetic turbulence, may even exceed the Spitzer value (Cho et al. (2003)) or be suppressed well below it. For our viscous simulations we choose f_v between 0 and 1 (see Table 1).

In order to ensure numerical stability and to prevent too small timesteps we have switched off viscosity for temperatures outside the range $3 \cdot 10^5 \text{ K} < T < 1.5 \cdot 10^8 \text{ K}$.

The total energy fluxes are not modified by viscosity as the effects of viscous heating are small.

2.1 Code

The simulations were performed with the FLASH code (Fryxell et al. 2000) version 2.5, a multidimensional adaptive mesh refinement hydrodynamics code. It solves the Riemann problem on a Cartesian grid using the Piecewise-Parabolic Method (PPM).

The viscous runs are fairly expensive because the

Table 1. ICM conditions for different simulation runs.

run:	M08	M08-V1	M08-V2	M08-V3
$\rho_{\text{ICM}}/10^{-27} \text{ g cm}^{-3}$	1	1	1	1
$v_{\text{ICM}}/1000 \text{ km s}^{-1}$	0.8	0.8	0.8	0.8
f_v	0	0.1	0.5	1.0

run:	M20	M20-V
$\rho_{\text{ICM}}/10^{-27} \text{ g cm}^{-3}$	1	1
$v_{\text{ICM}}/1000 \text{ km s}^{-1}$	2	2
f_v	0	0.1

timestep imposed by the viscosity scales as $(\Delta x)^2/\mu$, where Δx is the resolution of the computational grid. As this timestep scales more strongly with Δx than the standard hydrodynamical Courant condition, and because viscosity depends strongly on temperature, the constraints on the timestep are more stringent than in the inviscid runs. Hence, here we present only results from 2D simulations in cylindrical coordinates, (R, Z) .

The ICM wind enters the simulation box at $Z = -65 \text{ kpc}$. The boundary at $R = 0$ is the galaxy's symmetry axis, the remaining two boundaries obey outflow conditions. We use a grid size of 260 kpc in Z -direction 65 kpc in R -direction. With 5 refinement levels we reach an effective number of 2048×256 grid cells and effective resolution of 0.25 kpc.

2.2 Model galaxy

The galaxy model is the same as in Roediger & Brüggen (2006), i.e. a massive spiral with a flat rotation curve at 200 km s^{-1} . It consists of a dark matter halo ($1.1 \cdot 10^{11} M_\odot$ within 23 kpc), a stellar bulge ($10^{10} M_\odot$), a stellar disc ($10^{11} M_\odot$) and a gaseous disc ($5 \cdot 10^9 M_\odot$). All non-gaseous components just provide the galaxy's potential and are not evolved during the simulation. For a description of the individual components and a list of parameters please refer to RB06. Initially, the gas disc is set in hydrostatic equilibrium with the surrounding ICM (see also RB06). The disc's rotation is included via the centrifugal force.

2.3 ICM conditions

Table 1 lists the ICM conditions for the simulation runs. The ICM temperature was $7.2 \cdot 10^7 \text{ K}$ or 6.2 keV in all runs. We start the simulation with the ICM at rest and then increase the inflow velocity over the first 50 Myr from zero to v_{ICM} (for more details see RB06).

3 RESULTS

Figure 1 shows the density in the subsonic runs with increasing viscosity: $f_v = 0, 0.1$ and 0.5 . The different behaviour of the galactic wakes is evident: The larger the viscosity, the less turbulent the wake. In the viscous cases, the ICM flows past the galaxy more smoothly than in the non-viscous runs. In the non-viscous cases, the KH- and RT-instability cause vortices and turbulence. Both instabilities are suppressed

significantly in the viscous cases. Consequently, the stripped gas remains in larger clumps and is less readily mixed with the ambient medium.

Also the width of the tail, or the flaring angle, is slightly smaller in the viscous runs. In the inviscid run, the maximum width of the tail is $\approx 2 \times 40 \text{ kpc}$, while it is $\approx 2 \times 35 \text{ kpc}$ in all viscous runs. Thus, it is also unlikely that the small width of the tail in NGC 4388 and the X-ray tails of ESO137-001 (Sun et al. 2006) can be explained by a microphysical, Spitzer-type viscosity. Interestingly, the tail appears to be shorter, the larger the viscosity. This is due to the fact that larger gas clouds, even if they have the same density like smaller clouds, experience a smaller acceleration for a fixed ram pressure. We note that the mass fraction of stripped gas in dense form does not depend strongly on viscosity. In the viscous cases, there are a few large dense clouds, whereas in the non-viscous case there are numerous small dense clouds. We do not show any plots here for the supersonic runs but they show the same qualitative behaviour.

Figure 2 compares the evolution of the mass and radius of the remaining gas disc for viscous and inviscid runs. The top panel demonstrates that the viscosity has nearly no influence on the evolution of the gas disc's radius. The bottom panel displays the gas mass in a cylinder of radius 27 kpc and a height of 10 kpc. Interestingly, the mass evolution in all runs is very similar, for all values of the viscosity that we have explored. The first dip in the graph marks the end of the instantaneous stripping phase. Very slight differences introduced by the viscosity occur during the next, the intermediate phase and also during the final continuous stripping phase. We have also measured the gas mass bound to the gravitational potential of the galaxy and its behaviour is shown in the middle panel of Fig. 2. The differences between runs with and without viscosity is similar to that of the gas disc mass. This implies that the amount of gas lost from the galaxy is fixed by the ram pressure.

This result shows that the viscosity in the parameter range considered here has a minor impact on the mass loss history of the gas disc and that ram pressure pushing is the dominant mechanism, as found in earlier papers (e.g. Roediger & Brüggen 2006, 2007). The fact that the viscosity has a minor impact on the mass loss history should also hold for most galaxies which move through clusters and thus experience a variable ram pressure – for the very reason that ram pressure pushing is the main cause for mass loss also for these galaxies (Roediger & Brüggen 2007). The mass loss due to ram pressure pushing, i.e. during the first, instantaneous stripping phase, is especially insensitive to different ICM viscosities. An exception may occur when galaxies move near edge-on.

4 DISCUSSION

Clearly, we have to be careful with quantitative predictions especially concerning the morphology of the wake because our simulations are two-dimensional. However, in previous work we have shown that mass loss rates are the same in 2D and 3D simulations. Moreover, in the non-viscous cases, the wake structures in 2D and 3D were very similar. Thus, our main conclusions should be robust:




Figure 1. Contour plot of the density in subsonic runs at 300 Myr, for increasing viscosity from left to right: left is inviscid ($f_v = 0$), middle is viscous with $f_v = 0.1$, right is viscous with $f_v = 0.5$.

- The mass loss from the gas discs is hardly influenced by viscosity.
- With increasing viscosity, the wake shows less structure and turbulence, but larger clumps.

A detailed investigation of the fate of the stripped gas requires several additions to our simplified model: a 3D treatment to allow for other inclinations than face-on, prescriptions for heating, cooling and thermal conduction. These processes do not only influence the temperature in the stripped gas, but also determine in which wavebands the stripped gas will be observable. For the fate of the stripped gas, viscosity may make a difference as it may affect the thermal history of the stripped gas. Gradients of density and temperature are smeared out in the wake in the presence of viscosity. This will affect the efficiency of heat conduction and evaporation of clumps in the galactic wakes. It will also affect the efficiency with which material in the wake can form stars as observed by Sun et al. (2007). Comparisons between models and observations will reveal which processes are the dominant ones in shaping galactic tails.

ACKNOWLEDGEMENTS

We acknowledge the support by the DFG grant BR 2026/3 within the Priority Programme “Witnesses of Cosmic History” and the supercomputing grants NIC 2195 and 2256 at the John-Neumann Institut at the Forschungszentrum Jülich. The results presented were produced using the FLASH code, a product of the DOE ASC/Alliances-funded Center for Astrophysical Thermonuclear Flashes at the University of Chicago. We thank the referee for the helpful comments.

REFERENCES

- Abadi M. G., Moore B., Bower R. G., 1999, MNRAS, 308, 947
 Betchov R., Criminale W. O., 1967, Stability of parallel flows. Academic press, New York
 Braginskii S. L., 1958, Soviet Phys.-JETP Lett., 6, 358
 Cho J., Lazarian A., Vishniac E. T., 2003, ApJ, 595, 812
 Clarke T. E., 2004, JKAS, 37, 337
 Enßlin T. A., Vogt C., Pfrommer C., 2005, in Chyzy K. T.,




Figure 2. Evolution of radius (top panel) and mass of the remaining gas disc: the bottom panel measures the mass in a fixed cylinder centred on the galaxy, the middle panel measures the mass gravitationally bound to the galaxy. The viscous runs are for $f_v = 0.5$.

Otmianowska-Mazur K., Soida M., Dettmar R.-J., eds, The magnetized plasma in galaxy evolution Magnetic fields in clusters of galaxies. p. 231
 Fabian A. C., Sanders J. S., Allen S. W., Crawford C. S., Iwasawa K., Johnstone R. M., Schmidt R. W., Taylor G. B., 2003, MNRAS, 344, L43
 Fabian A. C., Sanders J. S., Crawford C. S., Conselice C. J., Gallagher J. S., Wyse R. F. G., 2003, MNRAS, 344, L48
 Fryxell B., Olson K., Ricker R., Timmes F. X., Zingale M., Lamb D. Q., MacNeice P., Rosner R., Truran J. W., Tufo H., 2000, ApJS, 131, 273
 Gunn J. E., Gott J. R., 1972, ApJ, 176, 1
 Marcolini A., Brighenti F., A.D'Ercole 2003, MNRAS, 345, 1329
 Narayan R., Medvedev M. V., 2001, ApJ, 562, L129
 Nulsen P. E. J., 1982, MNRAS, 198, 1007
 Oosterloo T., van Gorkom J., 2005, A&A, 437, L19
 Quilis V., Moore B., 2001, ApJ, 555, L95
 Quilis V., Moore B., Bower R., 2000, Science, 288, 1617
 Reynolds C. S., McKernan B., Fabian A. C., Stone J. M., Vernaleo J. C., 2005, MNRAS, 357, 242
 Roediger E., Brüggen M., 2006, MNRAS, 369, 567
 Roediger E., Brüggen M., 2007, MNRAS, 380, 1399
 Roediger E., Brüggen M., 2008, MNRAS, accepted

Roediger E., Brüggen M., Hoeft M., 2006, MNRAS, 371, 609
 Roediger E., Hensler G., 2005, A&A, 433, 875
 Ruszkowski M., Brueggen M., Begelman M. C., 2004a, ApJ, 615, 675
 Ruszkowski M., Brueggen M., Begelman M. C., 2004b, ApJ, 611, 158
 Schuecker P., Finoguenov A., Miniati F., Böhringer H., & U. G. Briel, 2004, A&A, 426, 387
 Schulz S., Struck C., 2001, MNRAS, 328, 185
 Spitzer L., 1962, Physics of Fully Ionized Gases. Interscience Publishers (2nd edition), New York
 Sun M., Donahue M., Voit G. M., 2007, ApJ, in press
 Sun M., Jones C., Forman W., Nulsen P. E. J., Donahue M., Voit G. M., 2006, ApJ, 637, L81
 Vollmer B., 2003, A&A, 398, 525
 Vollmer B., Balkowski C., Cayatte V., van Driel W., Huchtmeier W., 2004, A&A, 419, 35
 Vollmer B., Braine J., Balkowski C., Cayatte V., Duschl W. J., 2001, A&A, 374, 824
 Vollmer B., Cayatte V., Balkowski C., Duschl W. J., 2001, ApJ, 561, 708
 Vollmer B., Cayatte V., Boselli A., Balkowski C., Duschl W. J., 1999, A&A, 349, 411
 Vollmer B., Huchtmeier W., van Driel W., 2005, A&A, 439, 921
 Vollmer B., Marcelin M., Amram P., Balkowski C., Cayatte V., Garrido O., 2000, A&A, 364, 532
 Vollmer B., Soida M., Otmianowska-Mazur K., Kenney J. D. P., van Gorkom J. H., Beck R., 2006, A&A, 453, 883
 Yagi M., Komiyama Y., Yoshida M., Furusawa H., Kashikawa N., Koyama Y., Okamura S., 2007, ApJ, 660, 1209

This paper has been typeset from a $\text{\LaTeX}$ file prepared by the author.